

IASSNS-HEP-94/88

CCNY-HEP 94/9

October 1994

Many-body States and Operator Algebra for Exclusion Statistics

DIMITRA KARABALI

Institute for Advanced Study

Princeton, NJ 08540

V.P. NAIR

Physics Department

City College of the City University of New York

New York, New York 10031

Abstract

We discuss many-body states and the algebra of creation and annihilation operators for particles obeying exclusion statistics.

Recently Haldane introduced a variant of fractional statistics for which the notion of exclusion, a generalization of the Pauli principle, rather than exchange or braiding properties, is the prime characteristic^[1]. This exclusion statistics appears naturally in the fractional quantum Hall effect, spin- $\frac{1}{2}$ antiferromagnetic chains and the Calogero-Sutherland model^[1–5].

Exclusion principle implies that the number of available one-particle states should change with increasing occupation of a state and hence exclusion statistics can be characterized by the change Δd_i in available states as the occupation number is changed by ΔN_i , i.e., by

$$\Delta d_i = - \sum_j g_{ij} \Delta N_j \quad (1)$$

g_{ij} are the parameters characterizing the statistics. d_i is the number of one-particle states available to the N_i -th particle with quantum numbers i , holding the labels of the $(N_i - 1)$ particles fixed. Particles obeying exclusion statistics were called g -ons in ref.[6], a name we shall also use.

The thermodynamic distributions for particles obeying exclusion statistics can be set up generalizing the long familiar combinatorial calculation of entropy for bosons and fermions. A number of thermodynamic properties have been studied in this framework^[4–6]. A question which naturally arises in this context is whether one can go beyond the thermodynamic formulation, attempting a microscopic description by explicitly constructing the many-body Hilbert space and further introducing creation and annihilation operators and appropriate commutation rules.

In this paper we discuss an algebra of creation and annihilation operators which realizes some of the general features following from Eq.(1). The operator algebra is different from algebras discussed in connection with the Calogero-Sutherland system^[7]; further it seems to be different from proposals for anyon commutation rules^[8] as well as q -deformations of boson and fermion algebras, although there are many similarities^[9,10].

We start by considering g -ons of a single energy ω , i.e., only one species. Let $|n\rangle$

denote an n -body state of g -ons. (We consider them to be in a cavity of volume $V = L^3$ and fixed momentum of the form $\frac{2\pi}{L} \times \text{integer}$). n can be taken as the eigenvalue of some hermitian operator N . Further, we can assume $\langle n|m \rangle = \delta_{n,m}$.

We now introduce an annihilation operator a by

$$a|n\rangle = f_n|n-1\rangle \quad (2)$$

By orthonormality of states, $\langle n-1|a|n\rangle = f_n$, which leads, by the definition of the adjoint, to

$$a^\dagger|n\rangle = f_{n+1}^*|n+1\rangle \quad (3)$$

For a sensible annihilation operator we need $a|0\rangle = 0$, or $f_0 = 0$ and the one-particle state can be defined as $a^\dagger|0\rangle$, so that $f_1^* = 1$.

The number operator N , by definition, must obey the commutation rules

$$[N, a] = -a, \quad [N, a^\dagger] = a^\dagger \quad (4)$$

from which it follows that $[N, a^\dagger a] = [N, aa^\dagger] = 0$. The number operator N can thus be written, in general, as a function of $a^\dagger a$ or $aa^\dagger$.

From Eq.(1) we see that for $g = 1/m$, m being an integer, we lose a state every time the occupation number increases by m . We thus consider that in a microscopic description, Eq.(1) implies that the maximal occupancy of the state is m . In terms of operators this is equivalent to the condition $(a^\dagger)^{m+1} = (a)^{m+1} = 0$. A basis for the Hilbert space consists of the states $|0\rangle, |1\rangle, \dots, |m\rangle$. By definition the number operator N satisfies the polynomial relation $N(N-1)(N-2)\dots(N-m) = 0$. Further it can be written as an m -th order polynomial in $a^\dagger a$ (or $aa^\dagger$),

$$N = \sum_{k=1}^m \lambda_k (a^\dagger a)^k \quad (5)$$

Applying this on $|0\rangle, |1\rangle, |2\rangle$, etc., and using Eqs.(2) and (3) we get the equations

$$\sum_{k=1}^m |f_n|^{2k} \lambda_k = n, \quad n = 1, 2, \dots, m \quad (6)$$

We can now solve these equations for λ_k and hence obtain N as a function of $|f_l|^2$, $l = 2, \dots, m$. Similarly we can write down a “commutation rule” expressing the relation between $aa^\dagger$ and $a^\dagger a$,

$$aa^\dagger = 1 + \sum_{k=1}^m \sigma_k (a^\dagger a)^k \quad (7)$$

where σ_k ’s satisfy the equations

$$\sum_{k=1}^m |f_n|^{2k} \sigma_k = |f_{n+1}|^2 - 1 \quad (8)$$

The general solution to Eqs.(6) and (8) is given by

$$\begin{aligned} \lambda_k &= \sum_{l=1}^m \frac{l}{k!} \frac{1}{\det M} \left[\frac{\partial^k}{(\partial x_l)^k} \det M \right]_{x_l=0} \\ \sigma_k &= \sum_{l=1}^m \frac{(x_{l+1} - 1)}{k!} \frac{1}{\det M} \left[\frac{\partial^k}{(\partial x_l)^k} \det M \right]_{x_l=0} \end{aligned} \quad (9)$$

where

$$\det M = -(x_1 x_2 \dots x_m) \prod_{i < j} (x_i - x_j) \quad (10)$$

and $M_{lk} = (x_l)^k$. We substitute $x_l = |f_l|^2$ after evaluating the right hand side of Eq.(9).

For example, for $m = 2$ (“semions”) we find

$$\begin{aligned} aa^\dagger &= 1 + \frac{|f_2|^6 - |f_2|^4 + 1}{|f_2|^4 - |f_2|^2} a^\dagger a - \frac{|f_2|^4 - |f_2|^2 + 1}{|f_2|^4 - |f_2|^2} (a^\dagger a)^2 \\ &= 1 - (1 - |f_2|^2) a^\dagger a - \frac{1 + (|f_2|^2 - 1)|f_2|^2}{|f_2|^2} (a^\dagger)^2 a^2 \end{aligned} \quad (11)$$

$$\begin{aligned} N &= \frac{|f_2|^4 - 2}{|f_2|^4 - |f_2|^2} a^\dagger a + \frac{2 - |f_2|^2}{|f_2|^4 - |f_2|^2} (a^\dagger a)^2 \\ &= a^\dagger a + \frac{2 - |f_2|^2}{|f_2|^2} (a^\dagger)^2 a^2 \end{aligned} \quad (12)$$

We further have $(a^\dagger)^3 = 0$, $a^3 = 0$. (The case $m = 2$ is similar to the case considered in ref.[9].)

Eqs.(5-10) give an $(m-1)$ -parameter set of operator algebras, the parameters being $x_l = |f_l|^2$, $l = 2, \dots, m$. Specializations of this class of algebras can be obtained by

imposing relations among $|f_l|^2$. Such relations can be written in the form of a recursion rule

$$|f_{n+1}|^2 = K(n, g)|f_n|^2, \quad n = 1, \dots \quad (13)$$

the choice of the function K now parametrizing the possible algebras. Once a $K(n, g)$ is chosen, Eq.(13) can give, upto a phase, a recursive determination of f_n 's and hence a representation for a and $a^\dagger$. Some restrictions on $K(n, g)$ follow from general arguments. In order to obtain $f_{m+1} = 0$ from Eq.(13), consistent with the maximum- m occupancy for a state, $K(n, g)$ must vanish for $n = m = 1/g$. Thus we may expect a factor like $(1 - gn)$ to some positive power. Further we have the limiting cases $K(n, g) = \frac{1-n}{n}$ for $g = 1$ and $K(n, g) = \frac{1+n}{n}$ for $g = 0$, corresponding to fermions and bosons respectively. $|f_{n+1}|^2$ is essentially the probability of introducing an extra g -on into a state with occupation number n . For m very large and n/m small, the system is very nearly bosonic and the exclusion effect is insignificant; we expect $|f_{n+1}|^2 \sim (1 - gn) + n$, corresponding to a spontaneous emission term proportional to the available states $(1 - gn)$ and a stimulated emission term $\sim n$. Examples of possible choices for $K(n, g)$ are

$$K(n, g) = \frac{(1 - gn)[1 + (1 - g)n]}{n} \quad (14a)$$

$$K(n, g) = \frac{(1 - gn)^g[1 + (1 - g)n]^{1-g}}{n} \quad (14b)$$

There are of course many other choices possible. We expect $|f_{n+1}|^2/|f_n|^2$ to be proportional to the number of ways of introducing an extra g -on into the n -filled state, which is related to the combinatorial entropy per particle, say $S(n)$, as $\sim e^{S(n)}$. The combinatorial rule introduced in ref.[4] gives the entropy as

$$\begin{aligned} S &= \log \left[\frac{1 + (1 - g)n}{1 - gn} \right] + nS(n) \\ S(n) &= \log \left[\frac{(1 - gn)^g[1 + (1 - g)n]^{1-g}}{n} \right] \end{aligned} \quad (15)$$

Eq.(15) suggests the choice (14b). However we would like to emphasize that there is no compelling argument for such a choice. In particular the thermodynamic description of

ref. [4] cannot be derived from a microscopic theory in the standard way, given that there is no positive density matrix associated with it^[6]. We cannot therefore expect to recover Wu's thermal distribution from Eq.(14b). (Note that the maximum occupancy being m is sufficient to ensure that the thermodynamic distribution for the number density $\bar{n} \rightarrow m$ as the temperature goes to zero.) Nevertheless we shall analyze choice (14b), and its appropriate generalization in the case of many species, in some detail below because it is consistent with expected limiting behaviours and interpolates smoothly between bosons and fermions. Other choices will give qualitatively similar features.

A further restriction on $K(n, g)$ is derived from the fact that $|f|^2$, being probabilities, cannot be negative. A closer inspection of Eq.(14b) reveals that there are inconsistencies for some values of the parameter g . The inconsistency appears because, for some values of g , one can eventually encounter negative $|f|^2$. Our approach from now on will be that we shall use Eq.(14b) only for $g = 1/m$.

Using the recursion rule (14b) we find

$$f_1 = 1$$

$$|f_n|^2 = \prod_{k=1}^{n-1} \frac{(1 - \frac{k}{m})^{\frac{1}{m}} (1 + k(1 - \frac{1}{m}))^{1 - \frac{1}{m}}}{k}, \quad 2 \leq n \leq m \quad (16a)$$

$$= 0, \quad n \geq m + 1 \quad (16b)$$

As we mentioned earlier, this provides, upto a phase, a matrix realization of a and $a^\dagger$; $a_{mn} = \delta_{m,n-1} f_n$, $a_{mn}^\dagger = \delta_{m,n+1} f_{n+1}^*$.

We now turn to the case of many species corresponding to different values of energy or other quantum numbers. For two species of g -ons we introduce creation and annihilation operators $a_i^\dagger, a_i$, $i = 1, 2$ and $f_{n_1, n_2}^{(i)}$ with

$$a_1 |n_1, n_2\rangle = f_{n_1, n_2}^{(1)} |n_1 - 1, n_2\rangle,$$

$$a_2 |n_1, n_2\rangle = f_{n_1, n_2}^{(2)} |n_1, n_2 - 1\rangle, \quad (17)$$

where $f_{0, n_2}^{(1)} = f_{n_1, 0}^{(2)} = 0$ and $f_{1, 0}^{(1)} = f_{0, 1}^{(2)} = 1$. Eq.(3) can be generalized to the case of

two-species as

$$\begin{aligned} |f_{n_1+1, n_2}^{(1)}|^2 &= K^{(1)}(n_1, n_2, g) |f_{n_1, n_2}^{(1)}|^2, & n_1 > 0 \\ |f_{n_1, n_2+1}^{(2)}|^2 &= K^{(2)}(n_1, n_2, g) |f_{n_1, n_2}^{(2)}|^2, & n_2 > 0 \end{aligned} \quad (18)$$

A suitable generalization of Eq.(14b) is

$$\begin{aligned} K^{(i)}(n_1, n_2, g) &= (1 + w_i) \prod_k \left(\frac{w_k}{1 + w_k} \right)^{g_{ki}} \\ w_i n_i + g_{ij} n_j &= 1 \end{aligned} \quad (19)$$

The many-body states and the representation of a_i and $a_i^\dagger$ can be constructed by solving these recursion relations. We shall first address the case where g_{ij} is diagonal, i.e., no mutual statistics, and for the parameters $g_{ii} = 1/m$ we shall obtain the full operator algebra. We shall later discuss the case with mutual statistics and the somewhat complicated structure of the corresponding many-body Hilbert space.

In the absence of mutual statistics, the recursion rules (18) reduce to

$$\begin{aligned} |f_{n_1+1, n_2}^{(1)}|^2 &= w_1^{g_1} (1 + w_1)^{1-g_1} |f_{n_1, n_2}^{(1)}|^2 \\ |f_{n_1, n_2+1}^{(2)}|^2 &= w_2^{g_2} (1 + w_2)^{1-g_2} |f_{n_1, n_2}^{(2)}|^2 \end{aligned} \quad (20)$$

where $g_i = g_{ii}$ and $n_i w_i = 1 - g_i n_i$. For consistency reasons outlined earlier in the case of one species of g -ons, we shall consider special values of g_i , in particular the case $g_1 = g_2 = 1/m$. We again find that the maximal occupancy for each species is m ; i.e., $(a_i)^{m+1} = (a_i^\dagger)^{m+1} = 0$. Further, the recursion rules (20) are sufficient to show that the Hilbert space is a product of the Hilbert spaces for each species. These properties clearly generalize to arbitrary number of species.

The recursion rules (20) do not completely determine the functions $f_{n_1, n_2}^{(i)}$. Specifically, $|f_{1, n_2}^{(1)}|$ and $|f_{n_1, 1}^{(2)}|$ for $n_1, n_2 \neq 0$, as well as the phases of all the f 's, are undetermined. Thus Eq.(20) does not suffice to get all the matrix elements of a_i and $a_i^\dagger$. This is, of

course, no different from the situation with bosons or fermions; one needs commutation rules relating operators for different species in order to obtain a complete operator algebra and determine all the matrix elements.

The rationale for the choice of off-diagonal commutation rules has to come from the exclusion principle in the following way. Although not specified explicitly, we have been dealing all along with free g -ons. Thus the quantum number labelling the species is the spatial momentum p with a corresponding energy $\omega(p)$. The property of exclusion reads, for $g_i = 1/m$, $(a_p)^{m+1} = (a_p^\dagger)^{m+1} = 0$. Eventhough the p -diagonal representation is appropriate for free particles, we can equally well use the coordinate representation. Since the coordinates x also provide a complete set of quantum numbers for the particles, we expect the exclusion property $(a(x))^{m+1} = 0$, $(a^\dagger(x))^{m+1} = 0$ to hold, where $a(x) = \sum_p e^{ipx} a_p$ and $a^\dagger(x)$ is the adjoint of $a(x)$. Of course, one could also use $a(q) = \sum_q u_q(p) a_p$ in q -diagonal representation, q 's being any complete set of quantum numbers and $u_q(p)$ being the appropriate functions. Thus in general we expect linear combinations of a 's to obey the exclusion principle, i.e.,

$$\left(\sum_i \alpha_i a_i\right)^{m+1} = \left(\sum_i \alpha_i^* a_i^\dagger\right)^{m+1} = 0 \quad (21)$$

where α_i are arbitrary.

From the definition of the number operator $N_i|n_1, n_2, \dots\rangle = n_i|n_1, n_2, \dots\rangle$, we find that

$$[N_i, a_j] = [N_i, a_j^\dagger] = 0, \quad i \neq j \quad (22)$$

Eqs.(21,22) will be our main guide for obtaining the full operator algebra for the case of many species. However they are still not sufficient to determine the algebra completely, so we shall further assume that

$$a_i a_j = R_{ij} a_j a_i, \quad i \neq j \quad (23)$$

where R_{ij} is a c -number. Consistency of Eq.(23) requires $R_{ij} R_{ji} = 1$. Further, $a_i^\dagger a_j^\dagger = R_{ji}^* a_j^\dagger a_i^\dagger$. Since N_i is a function of $a_i^\dagger a_i$, we seek to satisfy Eq.(22) with the slightly stronger

requirement $[a_i^\dagger a_i, a_j] = [a_i^\dagger a_i, a_j^\dagger] = 0$, $i \neq j$. The general solution to this equation, using Eq.(23), is

$$\begin{aligned} a_i^\dagger a_j &= R_{ij}^{-1} a_j a_i^\dagger \\ a_i a_j^\dagger &= R_{ij}^* a_j^\dagger a_i, \quad i \neq j \end{aligned} \quad (24)$$

Consistency of these equations gives $R_{ij}^* R_{ij} = 1$. Thus R_{ij} is a phase, $R_{ij} = e^{i\theta_{ij}}$. We must now impose Eq.(21). For two species, we have

$$(\alpha_1 a_1 + \alpha_2 a_2)^{m+1} = (\alpha_1 a_1)^{m+1} + (\alpha_2 a_2)^{m+1} + \sum_{k=1}^m C(k, m, R_{12}^*) ((\alpha_1 a_1)^{m+1-k} (\alpha_2 a_2)^k) \quad (25)$$

where

$$C(k, m, R) = \frac{(1 - R^{m+1})(1 - R^m) \dots (1 - R^{m+2-k})}{(1 - R)(1 - R^2) \dots (1 - R^k)}, \quad 1 \leq k \leq m \quad (26)$$

In particular $C(1, m, R) = \frac{(1 - R^{m+1})}{(1 - R)}$. Since $(a_1)^{m+1} = (a_2)^{m+1} = 0$, Eq.(21) can be satisfied if all the $C(k, m, R_{12}^*)$ vanish. This determines R_{12} . The generic solution for arbitrary m is of the form $R_{12} = \exp(\pm 2\pi i / (m+1))$. There can be more solutions for special choices of m . All solutions are a subset of the $(m+1)$ -th roots of unity. For example, for $m = 2$, $R_{12} = e^{\pm 2\pi i / 3}$; $m = 3$, $R_{12} = e^{\pm 2\pi i / 4}$; $m = 4$, $R_{12} = e^{\pm 2\pi i / 5}$, $e^{\pm 4\pi i / 5}$. Since we want to treat all m 's on an equal footing, we are going to use the generic solution $R_{12} = \exp(\pm 2\pi i / (m+1))$. The commutation rule becomes $a_1 a_2 = e^{2\pi i / (m+1)} a_2 a_1$ or $a_1 a_2 = e^{-2\pi i / (m+1)} a_2 a_1$. Notice that the second choice is obtained from the first by a relabelling $a_1 \leftrightarrow a_2$. The basic solution will be $R_{12} = e^{2\pi i / (m+1)}$, up to such relabellings.

When we have more than two species, the exclusion condition (21) can be satisfied as follows. Write

$$a_1 a_3 = R_{13} a_3 a_1, \quad a_2 a_3 = R_{23} a_3 a_2 \quad (27)$$

For the choice $R_{23} = R_{13}$,

$$A a_3 = R_{13} a_3 A$$

where $A = \alpha_1 a_1 + \alpha_2 a_2$. The binary expansion (25) suffices to simplify $(A + \alpha_3 a_3)^{m+1} = 0$. We get the same solution as before, viz., $R_{13} = R_{12}$; the choice $R_{12} = R_{13} = R_{23}$ obviously generalizes inductively to arbitrary number of species.

Recapitulating, the commutation rules we have so far are, for $g_i = 1/m$,

$$a_i a_i^\dagger = 1 + \sum_{k=1}^m \sigma_k (a_i^\dagger a_i)^k \quad (28a)$$

$$a_i a_j = \exp\left(\frac{2\pi i}{m+1}\right) a_j a_i \quad i < j \quad (28b)$$

$$a_i^\dagger a_j^\dagger = \exp\left(\frac{2\pi i}{m+1}\right) a_j^\dagger a_i^\dagger \quad i < j \quad (28c)$$

$$a_i^\dagger a_j = \exp\left(-\frac{2\pi i}{m+1}\right) a_j a_i^\dagger \quad i < j \quad (28d)$$

We have arrived at Eqs.(28) by using the simplifying ansatz (23), with R_{ij} being a c -number. In order to write these commutation relations for different species we have to introduce an ordering, for example $i < j$. The indices i, j , etc., being momentum labels, ordering is naturally possible only in one spatial dimension. Since many of the physical situations where exclusion statistics might be relevant are effectively one-dimensional^[1–5], this may not be a drastic limitation. It is possible that more general structures can be constructed taking R_{ij} to be an operator. Notice also that once we have chosen an ordering of momenta we cannot relabel $a_1 \leftrightarrow a_2$, etc. The commutation rules with $R_{ij} \leftrightarrow R_{ij}^*$ are thus distinct.

We can now write down $f_{n_1, n_2}^{(i)}$ thus providing an explicit matrix realization for a_i and $a_i^\dagger$. Consider two species, $i = 1, 2$. From Eqs.(17,28) we derive two sets of equations that should be satisfied by f 's,

$$f_{n_1, n_2}^{(2)} f_{n_1, n_2-1}^{(1)} = \exp(2\pi i/(m+1)) f_{n_1, n_2}^{(1)} f_{n_1-1, n_2}^{(2)} \quad (29a)$$

$$f_{n_1, n_2}^{*(2)} f_{n_1, n_2}^{(1)} = \exp(-2\pi i/(m+1)) f_{n_1, n_2-1}^{(1)} f_{n_1-1, n_2}^{*(2)} \quad (29b)$$

Eqs. (20,29) completely determine the moduli $|f|$; in particular,

$$\begin{aligned} |f_{n_1, n_2}^{(1)}|^2 &= |f_{n_1, n_2-1}^{(1)}|^2 = \dots = |f_{n_1, 0}^{(1)}|^2 = |f_{n_1}|^2, & 0 \leq n_1, n_2 \leq m \\ |f_{n_1, n_2}^{(2)}|^2 &= |f_{n_1-1, n_2}^{(2)}|^2 = \dots = |f_{0, n_2}^{(2)}|^2 = |f_{n_2}|^2, & 0 \leq n_1, n_2 \leq m \end{aligned} \quad (30)$$

where $|f_{n_i}|^2$ are defined in Eq.(16) for a single species.

Eqs. (29) also determine some of the relative phases of the f 's, but they still allow some freedom of choice for the phases; this is related to the ambiguity in how many-particle states are defined. For example, $(a_1^\dagger)^{n_1}(a_2^\dagger)^{n_2}|0\rangle$ and $(a_2^\dagger)^{n_2}(a_1^\dagger)^{n_1}|0\rangle$ have the same number of particles but different phases. We will make the choice

$$|n_1, n_2, \dots\rangle = C (a_1^\dagger)^{n_1}(a_2^\dagger)^{n_2} \dots |0\rangle \quad (31)$$

where C is a normalization factor chosen to be real. This is ordered in terms of ascending momenta; the $a^\dagger$'s with smaller labels appear to the left. With this phase convention, we have, for two species and $m = 2$,

$$\begin{aligned} f_{1,0}^{(1)} &= f_{1,1}^{(1)} = f_{1,2}^{(1)} = f_{0,1}^{(2)} = 1 \\ f_{2,0}^{(1)} &= f_{2,1}^{(1)} = f_{2,2}^{(1)} = f_{0,2}^{(2)} = \left[\frac{\sqrt{3}}{2} \right]^{1/2} \\ f_{1,1}^{(2)} &= e^{2\pi i/3}, \quad f_{1,2}^{(2)} = e^{2\pi i/3} \left[\frac{\sqrt{3}}{2} \right]^{1/2} \\ f_{2,1}^{(2)} &= e^{4\pi i/3}, \quad f_{2,2}^{(2)} = e^{4\pi i/3} \left[\frac{\sqrt{3}}{2} \right]^{1/2} \end{aligned} \quad (32)$$

All other f 's are zero. We thus have a complete specification of a_i , $a_i^\dagger$. This can be easily extended to many species. In fact, it is evident that the commutation rules (28) suffice to evaluate the action of a_i 's and $a_i^\dagger$'s on any state of the form (31). The exchange properties of the many-body wavefunctions are evident from the representation (31) and the commutation rules (28).

In the case of mutual statistics, viz., g_{ki} not diagonal in Eq.(19), the structure of the many-body Hilbert space is somewhat more complicated. The number of many-body states can be greater or smaller, depending on the signs and magnitudes of g_{ki} , than what is expected from the tensor product of states for each species. This change in the expected number of states has to do with Eqs.(18,19). From Eq.(19), w_i can become negative

as n_j increases. Since $|f^{(i)}|^2$ is positive, a negative w_i can give an inconsistency, as we have already discussed earlier, unless some of the f 's are zero. Notice that Eq.(18) leaves $f_{1,n_2}^{(1)}, f_{n_1,1}^{(2)}$, for $n_1, n_2 \neq 0$ undetermined. One can set some of these to zero to obtain a consistent solution to Eqs.(18,19). The possible states are then no longer what is expected from the tensor product.

We shall illustrate this by considering two characteristic examples with two species, where the number of the states increases and decreases respectively, compared to the tensor product of the single species states.

A) $g_{11} = g_{22} = \frac{1}{m}, g_{12} = g_{21} = -\frac{1}{m}$

In this case the recursion rules reduce to

$$|f_{n_1+1,n_2}^{(1)}|^2 = (1 + w_1)^{1-\frac{1}{m}} \left(\frac{w_1}{w_2}\right)^{\frac{1}{m}} (1 + w_2)^{\frac{1}{m}} |f_{n_1,n_2}^{(1)}|^2 \quad (33a)$$

$$|f_{n_1,n_2+1}^{(2)}|^2 = (1 + w_2)^{1-\frac{1}{m}} \left(\frac{w_2}{w_1}\right)^{\frac{1}{m}} (1 + w_1)^{\frac{1}{m}} |f_{n_1,n_2}^{(2)}|^2 \quad (33b)$$

where $w_1 n_1 = 1 - \frac{(n_1 - n_2)}{m}$, $w_2 n_2 = 1 + \frac{(n_1 - n_2)}{m}$. We see that the ratio w_1/w_2 becomes negative if $|n_1 - n_2| > m$. A consistent solution requires that $f_{1,m+i}^{(1)} = 0$ and $f_{m+i,1}^{(2)} = 0$, $i \geq 1$. This choice allows only nonnegative w_1/w_2 . (This also avoids the singular points $w_2 = 0$ in Eq.(33a) and $w_1 = 0$ in Eq.(33b)). The allowed states are of the form $|n_1, n_2\rangle$, where $|n_1 - n_2| \leq m$. There are $(2m + 1)(m + 1)$ such states, which are $m(m + 1)$ more than in the tensor product of $(m + 1)$ states for each species.

B) $g_{11} = g_{22} = \frac{1}{m}, g_{12} = g_{21} = \frac{1}{m}$.

In this case the recursion rules become

$$|f_{n_1+1,n_2}^{(1)}|^2 = (1 + w_1)^{1-\frac{1}{m}} (w_1 w_2)^{\frac{1}{m}} (1 + w_2)^{-\frac{1}{m}} |f_{n_1,n_2}^{(1)}|^2 \quad (34a)$$

$$|f_{n_1,n_2+1}^{(2)}|^2 = (1 + w_2)^{1-\frac{1}{m}} (w_1 w_2)^{\frac{1}{m}} (1 + w_1)^{-\frac{1}{m}} |f_{n_1,n_2}^{(2)}|^2 \quad (34b)$$

where $w_1 n_1 = w_2 n_2 = 1 - \frac{(n_1 + n_2)}{m}$. Here we must require $f_{1,m+i}^{(1)} = 0$ and $f_{m+i,1}^{(2)} = 0$, $i \geq 0$, so as to avoid recursive sequences which lead to the singular points $w_2 = -1$ for Eq.(34a)

and $w_1 = -1$ for Eq.(34b). The allowed states now have $(n_1 + n_2) \leq m$. There are $\frac{1}{2}(m+2)(m+1)$ such states, which are $\frac{1}{2}m(m+1)$ less than in the tensor product of $(m+1)$ states for each species.

It is clear from the above discussion that the operator algebra for the case of mutual statistics must be fairly involved so as to reflect the complicated structure of the many-body states. This is currently under investigation.

Acknowledgements: The commutation rules (5,7) can be mapped into the usual bosonic algebra^[11]. We thank A.P. Polychronakos for discussions and for pointing this out.

This work was supported in part by the DOE grant DE-FG02-90ER40542 and the NSF grant PHY-9322591.

Note added: The combinatorial argument after Eq.(14), with equal a priori probability and maximal occupancy m , gives yet another choice for $K(n, g = 1/m)$ as

$$n = \frac{1}{K-1} - \frac{m+1}{K^{m+1}-1}$$

It is unclear how to generalize this equation to the case of mutual statistics. The case $m = 2$ for the above choice of K is similar to the case considered in ref.[12].

References

- [1] F.D.M. Haldane, *Phys. Rev. Lett.* **67** (1991) 937.
- [2] F.D.M. Haldane and B.S. Shastri, in the *Proceedings of the 16th Taniguchi Symposium*, Kashikojima, Japan, October 1993, eds. A. Okiji and N. Kawakami, Springer-Verlag, 1994.

- [3] Z.N.C. Ha, *Phys. Rev. Lett.* **73** (1994) 1574; preprint IASSNS-HEP-94/90, cond-mat/9410101; J. Minahan and A. Polychronakos, *Phys. Rev.* **B50** (1994) 4236; F. Lesage, V. Pasquier and D. Serban, preprint SPhT-94, hep-th/9405008.
- [4] Y.S. Wu, *Phys. Rev. Lett.* **73** (1994) 922.
- [5] D. Bernard and Y.S. Wu, preprint SPhT-94-043, UU-HEP/94-03, cond-mat/ 9404025; A. Dasnieres de Veigy and S. Ouvry, *Phys. Rev. Lett.* **72** (1994) 600; M.V.N. Murthy and R. Shankar, *Phys. Rev. Lett.* **72** (1994) 3629; preprint IMSc-94/24, cond-mat/9404096.
- [6] C. Nayak and F. Wilczek, preprint PUPT 1466, IASSNS 94/25, cond-mat/ 9405017, to appear in *Phys. Rev. Lett.*
- [7] A. Polychronakos, *Nucl. Phys.* **B324** (1989) 597; D.V. Khveshchenko, Princeton University preprint, cond-mat/9404094.
- [8] G. Semenoff, *Phys. Rev. Lett.* **61** (1988) 517; T. Matsuyama, *Phys. Lett.* **B228** (1989) 99; G. Semenoff and P. Sodano, *Nucl. Phys.* **B328** (1989) 753; D. Karabali, *Int. J. Mod. Phys.* **A6** (1991) 1369.
- [9] A.Y. Ignat'ev and V.A. Kuz'min, *Sov. J. Nucl. Phys.* **46** (1987) 786.
- [10] O.W. Greenberg and R.N. Mohapatra, *Phys. Rev. Lett.* **59** (1987) 2507; *Phys. Rev. Lett.* **62** (1989) 712; O.W. Greenberg, *Phys. Rev. Lett.* **64** (1990) 705; R.N. Mohapatra, *Phys. Lett.* **B242** (1990) 407.
- [11] A.P. Polychronakos, *Mod. Phys. Lett.* **A5** (1990) 2325.
- [12] P. Mitra, preprint SISSA-191/94/EP, hep-th/9411236.